

# Self doping effect and successive magnetic transitions in superconducting $\text{Sr}_2\text{VFeAsO}_3$

Guang-Han Cao,<sup>1,2,\*</sup> Zhifeng Ma,<sup>1</sup> Cao Wang,<sup>1</sup> Yunlei Sun,<sup>1</sup> Jinke Bao,<sup>1</sup> Shuai Jiang,<sup>1</sup> Yongkang Luo,<sup>1</sup> Chunmu Feng,<sup>1</sup> Yi Zhou,<sup>1</sup> Zhi Xie,<sup>3</sup> Fengchun Hu,<sup>3</sup> Shiqiang Wei,<sup>3</sup> I. Nowik,<sup>4</sup> I. Felner,<sup>4</sup> Lei Zhang,<sup>5</sup> Zhu'an Xu,<sup>1,2</sup> and Fu-Chun Zhang<sup>1,6</sup>

<sup>1</sup>Department of Physics, Zhejiang University, Hangzhou 310027, China

<sup>2</sup>State Key Lab of Silicon Materials, Zhejiang University, Hangzhou 310027, China

<sup>3</sup>National Synchrotron Radiation Laboratory, University of Science and Technology of China, Hefei 230029, China

<sup>4</sup>Racah Institute of Physics, The Hebrew University, Jerusalem 91904, Israel

<sup>5</sup>High Magnetic Field Laboratory, Chinese Academy of Sciences, Hefei 230031, China

<sup>6</sup>Department of Physics, The University of Hong Kong, Hong Kong, China

(Dated: September 28, 2010)

We have studied a quinary Fe-based superconductor  $\text{Sr}_2\text{VFeAsO}_3$  by the measurements of x-ray diffraction, x-ray absorption, Mössbauer spectrum, resistivity, magnetization and specific heat. This apparently undoped oxyarsenide is shown to be self doped via electron transfer from the  $\text{V}^{3+}$  ions. We observed successive magnetic transitions within the  $\text{VO}_2$  layers: an antiferromagnetic transition at 150 K followed by a weak ferromagnetic transition at 55 K. The spin orderings within the  $\text{VO}_2$  planes are discussed based on mixed valence of  $\text{V}^{3+}$  and  $\text{V}^{4+}$ .

PACS numbers: 74.70.Xa; 75.30.Kz; 71.28.+d; 74.10.+v

## I. INTRODUCTION

The discovery of high-temperature superconductivity (HTSC) in layered Fe-based compounds<sup>1,2</sup> represents an important breakthrough in the field of condensed matter physics. So far, dozens of new Fe-based superconductors in several crystallographic types have been discovered.<sup>2</sup> The common structural unit for the HTSC is  $\text{Fe}_2\text{X}_2$  ( $\text{X}=\text{As}, \text{Te}, \text{etc.}$ ) layers. The undoped  $\text{Fe}_2\text{X}_2$  layers with formally divalent iron usually exhibit antiferromagnetic (AFM) spin-density-wave (SDW) instability.<sup>3,4</sup> HTSC emerges as the SDW ordering is suppressed by various kinds of doping<sup>1,5–10</sup> or applying pressures<sup>11</sup>. Both the magnetic ordering and superconductivity were suggested to be related to the nesting between the two cylinder-like Fermi surfaces near  $\Gamma$  and M points.<sup>12–16</sup>

Recently, superconductivity at 37 K was reported in a quinary oxyarsenide  $\text{Sr}_2\text{VFeAsO}_3$  (hereafter called V21113),<sup>17</sup> consisting of perovskite-like  $\text{Sr}_4\text{V}_2\text{O}_6$  and anti-fluorite-type  $\text{Fe}_2\text{As}_2$  block layers. The observed superconductivity *without extrinsic doping* challenges the above paradigm, and immediately aroused several first principles calculations.<sup>18–22</sup> However, the occurrence of superconductivity in the apparently undoped V21113 has not been clarified. Moreover, the calculated property of the  $\text{Sr}_4\text{V}_2\text{O}_6$  layers is very much scattered: including magnetic half-metallic,<sup>18</sup> nonmagnetic metallic,<sup>19,20</sup> and magnetic (Mott-like) insulating<sup>22</sup> states. Various kinds of possibilities on the magnetic ground state for V21113 were evaluated, but no conclusive result was given.<sup>21</sup> Experimentally, few works<sup>17,23</sup> have been devoted to this new system. It is simply not clear whether the Fe-site AFM order survives, let alone the magnetic property of the  $\text{Sr}_4\text{V}_2\text{O}_6$  layers. The valence state of V, which is very crucial to understand the appearance of superconductivity, remains a puzzle. X-ray photoelectron spectroscopy (XPS) measurement suggested an "unex-

pected"  $\text{V}^{5+}$  even in the oxygen-deficient V21113 samples, however, the result was then questioned because of the surface-sensitive nature for the XPS method.<sup>23</sup>

In this article, we report systematic experimental studies on the structure and physical properties of V21113. We employed x-ray absorption spectroscopy, capable of detecting the bulk properties, to examine the valence state of V. Our data unambiguously reveal mixed valence of  $\text{V}^{3+}$  and  $\text{V}^{4+}$ , indicating that the  $\text{Fe}_2\text{As}_2$  layers are actually self doped due to the electron charge transfer from V to Fe. Our measurements have also identified two magnetic transitions at 55 K and 150 K, respectively, associated with the  $\text{VO}_2$  layers. The observed weak ferromagnetism below 55 K and its coexistence with superconductivity at low temperatures make the V21113 system very unique in the family of Fe-based superconductors.

## II. EXPERIMENTAL

Synthesis of single-phase 21113 samples turned out to be difficult,<sup>17</sup> because of the tendency of non-stoichiometry<sup>24</sup> as well as the multi-interphase solid-state reactions. We have improved the sample quality by carefully controlling the stoichiometry. Samples of  $\text{Sr}_2\text{V}_{1-x}\text{Mg}_x\text{FeAsO}_3$  ( $x=0$  corresponds to undoped V21113, and  $x=0.1$  refers to Mg-doped sample) were prepared using fine powders ( $\sim 200$  mesh, purity  $\geq 99.9\%$ ) of SrO, MgO, V, Fe,  $\text{Fe}_2\text{O}_3$  and As. The accurately-weighed stoichiometric mixture was loaded in an alumina tube, preventing possible reactions with the quartz tube. The alumina tube, placed in a sealed quartz ampoule, was heated slowly to 1273 K, holding for 24 h. After the first time sintering, the sample was thoroughly ground, pressed into pellets, and sintered again in vacuum ( $< 10^{-4}$  Pa) at 1353 K for 48 h. All the operations of weighing, mixing, grinding, pelletizing, etc. were car-

ried out in an argon-filled glove-box with the water and oxygen contents less than 0.1 ppm.

Several measurements were performed as follows. Powder x-ray diffraction (XRD) was carried out using a D/Max-rA diffractometer with Cu  $K_\alpha$  radiation and a graphite monochromator. The crystal structure at room temperature was refined by Rietveld analysis. The V  $K$ -edge x-ray absorption spectra (XAS) were measured in transmission mode using powdered samples on beam lines U7B and U7C at the National Synchrotron Radiation Facility (NSRF), Hefei, China. The water-cooled Si (111) plane bicrystal monochromator was used. Calibration of spectrometer was made using vanadium metal foil. The raw data were normalized according to the previous literature.<sup>25</sup> Mössbauer spectra were collected using a conventional constant acceleration drive. The sources were 50 mCi  $^{57}\text{Co}:\text{Rh}$ . The velocity calibration was performed with an  $\alpha$ -iron foil at room temperature. The reported isomer shift (IS) values for iron are relative to the Fe foil. The resistivity was measured on bar-shaped samples using a standard four-terminal method. We employed Quantum Design PPMS-9 and MPMS-5 to measure the specific heat and dc magnetization, respectively.

### III. RESULTS AND DISCUSSION

The XRD patterns are shown in Fig. 1(a). In comparison with the previous report<sup>17</sup>, the intensity of impurity peaks (relative to that of the main peak) at  $2\theta \sim 32^\circ$  is greatly reduced, indicating that the sample's quality is significantly improved. According to the multi-phase Rietveld refinement<sup>26</sup>, the amount of impurity phases  $\text{Sr}_2\text{VO}_4$  and FeAs are less than 5%. The refined structural parameters for  $\text{Sr}_2\text{V}_{1-x}\text{Mg}_x\text{FeAsO}_3$  ( $x=0$  and 0.1) are listed in Table I. The lattice constants of undoped V21113 are a little larger than the reported values<sup>17</sup>. The Mg substitution leads to a lattice elongation. The As–Fe–As angle is very close to the ideal value for a regular tetrahedron. On the other hand, the V–O2–V angle obviously deviates from  $180^\circ$ .

The low-temperature XRD experiments indicate no evidence of structural phase transition in the whole temperatures measured. Some of the data are shown in Fig. 1(b). First, the (200) peak does not separate within the experimental limit, suggesting no usual tetragonal-to-orthorhombic transition. Second, unlike the case in  $R\text{FeAsO}$  ( $R=\text{La}, \text{Sm}, \text{Gd}$  and  $\text{Tb}$ ),<sup>27,28</sup> no obvious splitting for the (220) peak can be seen. Thus, the SDW-ordering-related structural phase transition<sup>28</sup> can be ruled out in V21113. While the lattice parameters were found to decrease with decreasing temperature, the precise temperature dependence is expected to be measured by a future XRD study with synchrotron radiations.

The Mössbauer spectra [Fig. 1(c)] show a quadrupole doublet at all the temperatures, in contrast with the sextet line for the AFM SDW ordered state.<sup>27,29</sup> From




FIG. 1. (color online). (a) Rietveld refinement profiles for the x-ray diffraction (XRD) data of  $\text{Sr}_2\text{V}_{1-x}\text{Mg}_x\text{FeAsO}_3$  [ $x=0$  and 0.1 (inset)]. (b) Selected XRD peaks at various temperatures. (c)  $^{57}\text{Fe}$  Mössbauer spectra at different temperatures.

the data we deduced no effective paramagnetic moment, therefore, the iron does not carry magnetic moment on its own. Accordingly no magnetic ordering at the Fe sub-lattice is expected. The IS value is similar to those observed for other ferroarsenide systems,<sup>29</sup> suggesting that the iron is basically divalent.

Fig. 2 shows the V  $K$ -edge XAS for V21113 and its reference vanadium oxides. It has been well established that, in a number of V compounds, the position of the characteristic absorptions (such as absorption threshold, pre-edge peak, and main absorption edge) varies linearly with the valence of V ( $V_V$ ).<sup>25</sup> The main edge of V21113 lies in between those of  $\text{V}_2\text{O}_3$  and  $\text{VO}_2$ , clearly indicating the mixed valence state for V. The pre-edge structure (to the left of the main edge) of V21113 resembles that of  $\text{V}_2\text{O}_3$ , suggesting that  $\text{V}^{3+}$  is dominant. One can fit the pre-edge of V21113 by the addition of weighed XAS of  $\text{V}_2\text{O}_3$  and  $\text{VO}_2$ . The best fitting gives  $V_V=3.22$ . In fact,  $V_V$  can be better determined by using the characteristic absorption threshold ( $E_{\text{thres}}$ ). As shown in the insets of Fig. 2,  $E_{\text{thres}}$  increases steadily on going along  $\text{V}_2\text{O}_3 \rightarrow \text{VO}_2 \rightarrow \text{V}_2\text{O}_5$ . Assuming a linear relation of  $E_{\text{thres}}$  vs  $V_V$ , we obtain  $V_V = 3.18 \pm 0.05$ , consistent with the former estimation. Here we note that the structural parameters of V21113 also support the conclusion about the V valence. The calculated bond-valence-sum<sup>30</sup> values for V in Table I verify the dominant  $\text{V}^{3+}$  with minority of  $\text{V}^{4+}$  in V21113.

Because of the overall charge neutrality, the mixed va-

TABLE I. Crystallographic data of  $\text{Sr}_2\text{V}_{1-x}\text{Mg}_x\text{FeAsO}_3$  ( $x=0$  and 0.1) at room temperature. The space group is  $P4/nmm$ . The atomic coordinates are as follows: Sr1 (0.25, 0.25,  $z$ ); Sr2 (0.25, 0.25,  $z$ ); V (0.25, 0.25,  $z$ ); Fe (0.25, 0.75, 0); As (0.25, 0.25,  $z$ ); O1 (0.25, 0.25,  $z$ ); O2 (0.25, 0.75,  $z$ ). BVS refers to bond valence sum,<sup>30</sup> reflecting the formal valence of an ion investigated.

Compounds	$x=0$	$x=0.1$
$a$ (Å)	3.9352(1)	3.9334(2)
$c$ (Å)	15.6823(5)	15.7488(7)
$V$ (Å <sup>3</sup> )	242.86(1)	243.66(2)
$R_{wp}$	8.59	8.82
$S$	1.68	1.70
$z$ of Sr1	0.8090(1)	0.8091(1)
$z$ of Sr2	0.5857(1)	0.5865(1)
$z$ of V	0.3080(2)	0.3074(2)
$z$ of As	0.0893(1)	0.0880(2)
$z$ of O1	0.4248(4)	0.4257(5)
$z$ of O2	0.2934(3)	0.2952(4)
BVS <sub>V</sub> for V <sup>III</sup>	2.95	N.V.
BVS <sub>V</sub> for V <sup>IV</sup>	3.29	N.V.
BVS <sub>V</sub> for V <sup>V</sup>	3.46	N.V.
V–O2–V angle (°)	166.7(2)	168.8(2)
As–Fe–As angle (°)	109.1(1)	109.7(1)




FIG. 2. (color online). Vanadium  $K$ -edge x-ray absorption spectra for  $\text{Sr}_2\text{VFeAsO}_3$  as well as some reference compounds  $\text{V}_2\text{O}_3$ ,  $\text{VO}_2$  and  $\text{V}_2\text{O}_5$ . The insets show the determination of the valence of V ( $V_V$ ) by the absorption threshold,  $E_{\text{thres}}$ .

lence of V immediately suggests that the  $\text{Fe}_2\text{As}_2$  layers are "self" (or internally) electron-doped by the charge transfer from the  $\text{V}^{3+}$  ions (conversely, the  $\text{Sr}_4\text{V}_2\text{O}_6$  layers are self hole-doped). This conclusion is consistent with the measurements of Hall and Seebeck coefficients, which show electron-dominant transport behavior (not shown here), and also with the band calculations<sup>19</sup> which indicate the occupation number of  $3d$  electrons per V being less than 2. The absence of Fe-site AFM ordering and appearance of SC transition (see below) are thus naturally understood within the established paradigm for Fe-based HTSC.

In Fig. 3, we plot the temperature dependence of re-




FIG. 3. (color online). Temperature dependence of resistivity (a), magnetic susceptibility (b), and specific heat (c) for  $\text{Sr}_2\text{VFeAsO}_3$  and  $\text{Sr}_2\text{V}_{0.9}\text{Mg}_{0.1}\text{FeAsO}_3$ . The left axes show the results after different data processing.  $\chi_0$  is  $1.4 \times 10^{-3}$  and  $2.0 \times 10^{-3}$  emu/mol for the undoped and Mg-doped samples, respectively.

sistivity ( $\rho$ ), magnetic susceptibility ( $\chi$ ) and specific heat ( $C$ ) for V21113 along with  $\text{Sr}_2\text{V}_{0.9}\text{Mg}_{0.1}\text{FeAsO}_3$  as a reference. The V21113 sample shows superconducting (SC) transition at 24 K, which is somewhat lower than previously reported<sup>17</sup>. The lowered  $T_c$  is mainly ascribed to the "overdoped" state, since the self-doping level achieves  $\sim 18\%$  (from the obtained  $V_V$  above). As we see, the Mg-doped sample shows an enhanced  $T_c$  of 30 K. This result does suggest that the undoped V21113 be overdoped, because the Mg-for-V substitution induces holes, which partially compensates the heavy electron doping. Another reason for the lowered  $T_c$  could be due to the V-for-Fe anti-site occupation, as suggested by a very recent report.<sup>31</sup>

For V21113, an anomaly at  $T_1 = 150$  K is clearly shown in the derivative of  $\rho(T)$  as well as  $\chi(T)$  and  $C(T)$  curves. In particular, the anomaly in  $C(T)$  points to an intrinsic second-order phase transition, since specific heat measures bulk properties (note that the amount of impurities is less than 5%). For the Mg-doped sample, however, such an anomaly is completely suppressed. Since the Fe-site magnetic ordering was ruled out by the above Mössbauer study, we conclude that the 150-K transition V21113 is intrinsic and directly related to the V atoms.

The  $\chi(T)$  data in the range of  $300 \text{ K} > T > 150 \text{ K}$  obey the Curie-Weiss law,  $\chi = \chi_0 + C_{\text{Curie}}/(T - \theta_W)$ , where

$C_{\text{Curie}}$  denotes the Curie constant and  $\theta_W$  the Weiss temperature. Since the Fe atoms have no effective paramagnetic moments, as revealed by the above Mössbauer study, we can estimate the effective magnetic moments of V, which gives  $P_{\text{eff}}=1.2 \mu_B/\text{V}$  by the data fitting. The significantly lowered  $P_{\text{eff}}$  (compared with the spin-only value for either  $\text{V}^{3+}$  or  $\text{V}^{4+}$ ) could be due to the  $d-p$  hybridization in connection with the mixed valence. The fitted  $\theta_W$  value is 52 K, implying dominant ferromagnetic (FM) correlations between the magnetic moments. By subtracting  $C(T)$  of Mg-doped  $\text{V}_{21113}$  from that of the undoped  $\text{V}_{21113}$ , the magnetic entropy for the transition is calculated to be  $\sim 1.1 \text{ J}\cdot\text{mol}^{-1}\cdot\text{K}^{-1}$ , which is much lower than the expected value of  $R\ln(2S+1)$  ( $S=1$  for  $\text{V}^{3+}$ ; and  $S=1/2$  for  $\text{V}^{4+}$ ). The result suggests that short-range magnetic order is established at the temperatures far above 150 K. Note that the step-like increase in  $\chi$  at the transition does not mean a FM transition, because the  $M(H)$  curve for  $70 \text{ K} < T < 150 \text{ K}$  is linear [see Fig. 4(c)]. In fact, the AFM ordering of  $A$  or  $C$  type possibly exhibits an increase in  $\chi$  at the magnetic transition.<sup>32</sup>

In addition to the 150-K transition, another anomaly at  $T_2 \sim 55 \text{ K}$  can be clearly seen in  $1/(\chi - \chi_0)$ , as shown in Fig. 3(b). Specific heat data also show a small anomaly around 55 K. However, the Mg-doped sample, which contains a little more impurities, *simultaneously* loses the two anomalies. This fact strongly suggests that the 55-K anomaly is also intrinsic, and related to the  $\text{Sr}_4\text{V}_2\text{O}_6$  layers. Fig. 4 supplies more information about this second transition. At low fields, the magnetization diverges at 55 K for field-cooling (FC) and zero-field-cooling (ZFC) modes, suggesting a ferromagnetic transition. The FM state is further supported by the  $M(H)$  curve, where a magnetic hysteresis loop is clearly observed. The very small extrapolated residual magnetization ( $M_0 \sim 10^{-3} \mu_B/\text{V}$  at 35 K) and lack of saturation in  $M$  at high fields indicate that it is actually a weak ferromagnetism (WFM). The WFM nature is consistent with the above Mössbauer spectra which show negligibly small transferred hyperfine field (below 0.07 T) at the Fe site for  $T=20$  and 40 K. Nevertheless, the SC magnetic loop [Fig. 4(d)] exhibits obvious asymmetry relative to the horizontal axis, suggesting significant contributions of WFM in the SC state. It is noted that the coexistence of SC and WFM differs with the previous finding of the coexistence of SC and strong ferromagnetism (due to 4f local moments) in  $\text{EuFe}_2(\text{As}_{0.7}\text{P}_{0.3})_2$ .<sup>10</sup>

Now, let us discuss the possible origin of the two magnetic transitions within the  $\text{Sr}_4\text{V}_2\text{O}_6$  layers. The electronic configurations of  $\text{V}^{3+}$  and  $\text{V}^{4+}$  are  $d^2$  and  $d^1$ , respectively. Within an ionic model, the lowest V 3d orbitals are degenerate  $d_{yz}$  and  $d_{xz}$  due to the crystal field splitting. In the case of pure  $\text{V}^{3+}$ , the electron occupation is  $d_{yz}^\uparrow d_{xz}^\uparrow$  according to the Hund's rule. The ground state of the square lattice of V in the  $\text{VO}_2$  planes is AFM due to superexchange interaction.<sup>32</sup> In  $\text{V}_{21113}$ , there are  $\sim 18\%$   $\text{V}^{4+}$  ions, and the Zener's double exchange favors




FIG. 4. (color online). Low-field  $\chi(T)$  curves (a) and low-field  $M(H)$  curves (b) for  $\text{Sr}_2\text{VFeAsO}_3$  (in comparison with the Mg-doped sample). Panel (c) shows a full-range  $M(H)$  curves. Panel (d) displays the superconducting magnetization curve after subtracting a paramagnetic background.

ferromagnetic spin arrangement in the same  $\text{VO}_2$  plane for the gain of kinetic energy. This could result in an in-plane FM state. However, the electron hopping integrals in  $\text{VO}_2$  planes are direction-dependent. The amplitude of the hopping integral of  $d_{xz}$  along  $y$ -axis (or  $d_{yz}$  along  $x$ -axis),  $t'$ , is much smaller than that of  $d_{xz}$  along  $x$ -axis (or  $d_{yz}$  along  $y$ -axis),  $t$ . This leads to another possible spin array in the  $\text{VO}_2$  planes, i.e. a  $C$ -type AFM state where the spins are parallel along  $x$ -axis, and antiparallel along  $y$ -axis, or vice versa. By a simplified mean field analysis, we estimate that the FM state is more stable if  $|t'|/J > 2(1-\delta)/\delta$ , with  $J$  the neighboring spin coupling and  $\delta$  the concentration of  $\text{V}^{4+}$ .

The scenario of in-plane FM state coincides with the afore-mentioned positive Weiss temperature. Thus the observed transition at  $T_1=150 \text{ K}$  may be ascribed to an  $A$ -type AFM ordering with AFM coupling along the  $c$ -axis. The WFM transition at  $T_2=55 \text{ K}$  may be interpreted as the spin canting within the framework of double exchange proposed by de Gennes,<sup>33</sup> although usually one would expect a more substantial magnetic moment. As for the scenario of in-plane  $C$ -type AFM, the WFM transition is still an open question. One possibility is due to the nonzero Dzyaloshinsky-Moriya interaction, which plays an important role for the intriguing magnetic responses in other vanadium oxides.<sup>34</sup>

#### IV. CONCLUDING REMARKS

To summarize, we have demonstrated experimentally that superconductivity in undoped  $\text{Sr}_2\text{VFeAsO}_3$  is induced by a novel self doping mechanism, i.e. an inter-layer charge transfer from V to Fe. Neither structural phase transition nor Fe-site SDW ordering was observed,

consistent with the electron-doped state in the  $\text{Fe}_2\text{As}_2$  layers. In addition, we have identified two intrinsic magnetic transitions in connection with the mixed valence of V. The transition at 150 K is ascribed as an AFM ordering. The following transition at 55 K is weakly ferromagnetic, probably due to a spin canting process. The WFM transition renders a rare example of coexistence of SC and WFM, which will be of great interest for the future studies. Further experiments such as neutron diffractions are desirable to check the spin structure of V, and to confirm that the moment doesn't come from Fe.

## ACKNOWLEDGMENTS

This work is supported by the NSF of China (No. 90922002), National Basic Research Program of China (No. 2007CB925001) and the Fundamental Research Funds for the Central Universities of China (No. 2010QNA3026). The research in Jerusalem is supported by the Israel Science Foundation (Bikura 459/09) and by the joint German-Israeli DIP project. FCZ acknowledges partial support from Hong Kong RGC grant HKU 7068/09P and NSF/RGC N-HKU 726/09. The authors would like to thank NSRL for the synchrotron radiation beam time.

- 
- \* corresponding author: ghcao@zju.edu.cn
- <sup>1</sup> Y. Kamihara, T. Watanabe, M. Hirano, and H. Hosono, *J. Am. Chem. Soc.* **130**, 3296 (2008).
  - <sup>2</sup> For a recent review, see D. C. Johnston, arXiv:1005.4392 (2010).
  - <sup>3</sup> C. de la Cruz, Q. Huang, J. W. Lynn, J. Li, W. Ratcliff II, J. L. Zarestky, H. A. Mook, G. F. Chen, J. L. Luo, N. L. Wang, and P. Dai, *Nature (London)* **453**, 899 (2008).
  - <sup>4</sup> Q. Huang, Y. Qiu, W. Bao, M. A. Green, J. W. Lynn, Y. C. Gasparovic, T. Wu, G. Wu, and X. H. Chen, *Phys. Rev. Lett.* **101**, 257003 (2008).
  - <sup>5</sup> Z. A. Ren, J. Yang, W. Lu, W. Yi, X. L. Shen, Z. C. Li, G. C. Che, X. L. Dong, L. L. Sun, F. Zhou, and Z. X. Zhao, *EPL* **82**, 57002 (2008).
  - <sup>6</sup> H. H. Wen, G. Mu, L. Fang, H. Yang, and X. Zhu, *EPL* **82**, 17009 (2008).
  - <sup>7</sup> C. Wang, L. Li, S. Chi, Z. Zhu, Z. Ren, Y. Li, Y. Wang, X. Lin, Y. Luo, S. Jiang, X. Xu, G. Cao, and Z. Xu, *EPL* **83**, 67006 (2008).
  - <sup>8</sup> A. S. Sefat, A. Huq, M. A. McGuire, R. Y. Jin, B. C. Sales, D. Mandrus, L. M. D. Cranswick, P. W. Stephens, and K. H. Stone, *Phys. Rev. B* **78**, 104505 (2008); C. Wang, Y. K. Li, Z. W. Zhu, S. Jiang, X. Lin, Y. K. Luo, S. Chi, L. J. Li, Z. Ren, M. He, H. Chen, Y. T. Wang, Q. Tao, G. H. Cao, and Z. A. Xu, *Phys. Rev. B* **79**, 054521 (2009).
  - <sup>9</sup> G. H. Cao, S. Jiang, X. Lin, C. Wang, Y. K. Li, Z. Ren, Q. Tao, C. M. Feng, J. H. Dai, Z. A. Xu, and F. C. Zhang, *Phys. Rev. B* **79**, 174505 (2009).
  - <sup>10</sup> Z. Ren, Q. Tao, S. Jiang, C. M. Feng, C. Wang, J. H. Dai, G. H. Cao, and Z. A. Xu, *Phys. Rev. Lett.* **102**, 137002 (2009).
  - <sup>11</sup> M. S. Torikachvili, S. L. Budko, N. Ni, and P. C. Canfield, *Phys. Rev. Lett.* **101**, 057006 (2008).
  - <sup>12</sup> D. J. Singh and M.-H. Du, *Phys. Rev. Lett.* **100**, 237003 (2008).
  - <sup>13</sup> J. Dong, H. J. Zhang, G. Xu, Z. Li, G. Li, W. Z. Hu, D. Wu, G. F. Chen, X. Dai, J. L. Luo, Z. Fang, and N. L. Wang, *EPL* **83**, 27006 (2008).
  - <sup>14</sup> I. I. Mazin, D. J. Singh, M. D. Johannes, and M. H. Du, *Phys. Rev. Lett.* **101**, 057003 (2008).
  - <sup>15</sup> K. Kuroki, S. Onari, R. Arita, H. Usui, Y. Tanaka, H. Kon-tani, and H. Aoki, *Phys. Rev. Lett.* **101**, 087004 (2008).
  - <sup>16</sup> K. Terashima, Y. Sekiba, J. H. Bowen, K. Nakayama, T. Kawahara, T. Sato, P. Richard, Y. M. Xu, L. J. Li, G. H. Cao, Z. A. Xu, H. Ding, and T. Takahashi, *Proc. Nat. Acad. Sci. USA* **106**, 7330 (2009).
  - <sup>17</sup> X. Y. Zhu, F. Han, G. Mu, P. Cheng, B. Shen, B. Zeng, and H. H. Wen, *Phys. Rev. B* **79**, 220512(R) (2009).
  - <sup>18</sup> I. R. Shein and A. L. Ivanovskii, *J. Supercond. Nov. Magn.* **22**, 613 (2009).
  - <sup>19</sup> K. W. Lee and W. E. Pickett, *EPL* **89**, 57008 (2010).
  - <sup>20</sup> G. Wang, M. Zhang, L. Zheng, and Z. Yang, *Phys. Rev. B* **80**, 184501 (2009).
  - <sup>21</sup> I. I. Mazin, *Phys. Rev. B* **81**, 020507(R) (2010).
  - <sup>22</sup> H. Nakamura and M. Machida, arXiv:1004.4741 (2010).
  - <sup>23</sup> F. Han, X. Y. Zhu, G. Mu, P. Cheng, B. Shen, B. Zeng, and H. H. Wen, *Sci. China G* **53**, 1202 (2010).
  - <sup>24</sup> M. Tegel, F. Hummel, Y. X. Su, T. Chatterji, M. Brunelli, and D. Jorhendt, *EPL* **89**, 37006 (2010).
  - <sup>25</sup> J. Wong, F. W. Lytle, R. P. Messmer and D. H. Maylotte, *Phys. Rev. B* **30**, 5596 (1984).
  - <sup>26</sup> The reliable factors of the refinement are  $R_{wp}=0.086$  and  $R_{exp}=0.051$ , indicating a fairly good accuracy for the structural parameters obtained.
  - <sup>27</sup> M. A. McGuire, A. D. Christianson, A. S. Sefat, B. C. Sales, M. D. Lumsden, R. Y. Jin, E. A. Payzant, D. Mandrus, Y. B. Luan, V. Keppens, V. Varadarajan, J. W. Brill, R. P. Hermann, M. T. Sougrati, F. Grandjean, and G. J. Long, *Phys. Rev. B* **78**, 094517 (2008).
  - <sup>28</sup> Y. K. Luo, Q. Tao, Y. K. Li, X. Lin, L. J. Li, G. H. Cao, Z. A. Xu, Y. Xue, H. Kaneko, A. V. Savinkov, H. Suzuki, C. Fang, and J. P. Hu, *Phys. Rev. B* **80**, 224511 (2009).
  - <sup>29</sup> I. Nowik and I. Felner, *Physica C* **469**, 485 (2009).
  - <sup>30</sup> I. D. Brown and D. Altermatt, *Acta Cryst.* **B41**, 244 (1985).
  - <sup>31</sup> M. Tegel, T. Schmid, T. Stürzer, M. Egawa, Y. X. Su, A. Senyshyn, and D. Jorhendt, arXiv: 1008.2687.
  - <sup>32</sup> J. S. Zhou, J. B. Goodenough, J. Q. Yan, and Y. Ren, *Phys. Rev. Lett.* **99**, 156401 (2007).
  - <sup>33</sup> P. -G. de Gennes, *Phys. Rev.* **118**, 141 (1960).
  - <sup>34</sup> Y. Ren, T. T. M. Palstra, D. I. Khomskii, E. Pellegrin, A. A. Nugroho, A. A. Menovsky, and G. A. Sawatzky, *Nature* **396**, 441 (1998).